arXiv:1001.0326v1 [math.RT] 3 Jan 2010

Herz–Schur Multipliers and Non-Uniformly Bounded Representations of Locally Compact Groups

Troels Steenstrup*

Abstract

Let G be a second countable, locally compact group and let φ be a continuous Herz–Schur multiplier on G. Our main result gives the existence of a (not necessarily uniformly bounded) strongly continuous representation π of G on a Hilbert space $\mathscr{H}$, together with vectors $\xi, \eta \in \mathscr{H}$, such that $\varphi(y^{-1}x) = \langle \pi(x)\xi, \pi(y^{-1})^*\eta \rangle$ for $x, y \in G$ and $\sup_{x\in G} \|\pi(x)\xi\| \cdot \sup_{y\in G} \|\pi(y^{-1})^*\eta\| = \|\varphi\|_{M_0A(G)}$. Moreover, we obtain control over the growth of the representation in the sense that $\|\pi(g)\| \leq \exp(\frac{c}{2}\mathrm{d}(g, e))$ for $g \in G$, where $e \in G$ is the identity element, c is a constant and d is a metric on G.

Introduction

Let Y be a non-empty set. A function $\psi : Y \times Y \to \mathbb{C}$ is called a *Schur multiplier* if for every operator $A = (a_{x,y})_{x,y\in Y} \in \mathbf{B}(\ell^2(Y))$ the matrix $(\psi(x, y)a_{x,y})_{x,y\in Y}$ again represents an operator from $\mathbf{B}(\ell^2(Y))$ (this operator is denoted by $M_\psi A$). If ψ is a Schur multiplier it follows easily from the closed graph theorem that $M_\psi \in \mathbf{B}(\mathbf{B}(\ell^2(Y)))$, and one refers to $\|M_\psi\|$ as the *Schur norm* of ψ and denotes it by $\|\psi\|_S$.

Let G be a locally compact group. In [Her74], Herz introduced a class of functions on G, which was later denoted the class of *Herz–Schur multipliers* on G. By the introduction to [BF84], a continuous function $\varphi : G \to \mathbb{C}$ is a Herz–Schur multiplier if and only if the function

$$\hat{\varphi}(x, y) = \varphi(y^{-1}x) \qquad (x, y \in G) \tag{0.1}$$

---

*Partially supported by the Ph.D.-school OP–ALG–TOP–GEO.

is a Schur multiplier, and the *Herz–Schur norm* of φ is given by

$$\|\varphi\|_{HS} = \|\hat{\varphi}\|_S.$$

In [DCH85] De Cannière and Haagerup introduced the Banach algebra $MA(G)$ of *Fourier multipliers* of G, consisting of functions $\varphi : G \to \mathbb{C}$ such that

$$\varphi\psi \in A(G) \qquad (\psi \in A(G)),$$

where $A(G)$ is the *Fourier algebra* of G as introduced by Eymard in [Eym64] (the *Fourier–Stieltjes algebra* $B(G)$ of G is also introduced in this paper). The norm of φ (denoted $\|\varphi\|_{MA(G)}$) is given by considering φ as an operator on $A(G)$. According to [DCH85, Proposition 1.2] a Fourier multiplier of G can also be characterized as a continuous function $\varphi : G \to \mathbb{C}$ such that

$$\lambda(g) \overset{M_\varphi}{\mapsto} \varphi(g)\lambda(g) \qquad (g \in G)$$

extends to a σ-weakly continuous operator (still denoted M_φ) on the group von Neumann algebra ($\lambda : G \to \mathbf{B}(L^2(G))$ is the *left regular representation* and the group von Neumann algebra is the closure of the span of $\lambda(G)$ in the weak operator topology). Moreover, one has $\|\varphi\|_{MA(G)} = \|M_\varphi\|$. The Banach algebra $M_0A(G)$ of *completely bounded Fourier multipliers* of G consists of the Fourier multipliers of G, φ, for which M_φ is completely bounded. In this case they put $\|\varphi\|_{M_0A(G)} = \|M_\varphi\|_{\mathrm{cb}}$.

In [BF84] Bożejko and Fendler show that the completely bounded Fourier multipliers coincide isometrically with the continuous Herz–Schur multipliers. In [Jol92] Jolissaint gives a short and self-contained proof of the result from [BF84] in the form stated below.

0.1 Proposition ([BF84], [Jol92])**.** *Let G be a locally compact group and assume that $\varphi : G \to \mathbb{C}$ and $k \geq 0$ are given, then the following are equivalent:*

(i) *φ is a completely bounded Fourier multiplier of G with $\|\varphi\|_{M_0A(G)} \leq k$.*

(ii) *φ is a continuous Herz–Schur multiplier on G with $\|\varphi\|_{HS} \leq k$.*

(iii) *There exists a Hilbert space $\mathscr{H}$ and two bounded, continuous maps $P, Q : G \to \mathscr{H}$ such that*

$$\varphi(y^{-1}x) = \langle P(x), Q(y)\rangle \qquad (x, y \in G)$$

and

$$\|P\|_\infty \|Q\|_\infty \leq k,$$

where

$$\|P\|_\infty = \sup_{x\in G} \|P(x)\| \quad \text{and} \quad \|Q\|_\infty = \sup_{y\in G} \|Q(y)\|.$$

By a *representation* $(\pi, \mathscr{H})$ of a locally compact group G on a Hilbert space $\mathscr{H}$ we mean a homomorphism of G into the invertible elements of $\mathbf{B}(G)$. A representation $(\pi, \mathscr{H})$ of G is said to be *uniformly bounded* if

$$\sup_{g \in G} \|\pi(g)\| < \infty$$

and one usually writes $\|\pi\|$ for $\sup_{g \in G} \|\pi(g)\|$. If $g \mapsto \pi(g)$ is continuous with respect to the strong operator topology on $\mathbf{B}(G)$ then we say that $(\pi, \mathscr{H})$ is *strongly continuous*. Let $(\pi, \mathscr{H})$ be a strongly continuous, uniformly bounded representation of G then, according to [DCH85, Theorem 2.2], any *coefficient of* $(\pi, \mathscr{H})$ is a continuous Herz–Schur multiplier, i.e.,

$$g \overset{\varphi}{\mapsto} \langle \pi(g)\xi, \eta \rangle \qquad (g \in G)$$

is a continuous Herz–Schur multiplier with

$$\|\varphi\|_{M_0A(G)} \leq \|\pi\|^2 \|\xi\| \|\eta\|$$

for any $\xi, \eta \in \mathscr{H}$ (note that this result also follows as a corollary to Proposition 0.1). U. Haagerup has shown that on the non-abelian free groups there are Herz–Schur multipliers which can not be realized as coefficients of uniformly bounded representations. The proof by Haagerup has remained unpublished, but Pisier has later given a different proof, cf. [Pis05]. Haagerup's proof can be modified to prove the corresponding result for the connected, real rank one, simple Lie groups with finite center, cf. [Ste09, Theorem 3.6].

Strictly speaking, the requirement that the above representations be uniformly bounded is not fully needed in order to construct a continuous Herz–Schur multiplier. From Proposition 0.1 it follows that it is enough to require that

$$\sup_{x \in G} \|\pi(x)\xi\| < \infty \quad \text{and} \quad \sup_{y \in G} \|\pi(y^{-1})^*\eta\| < \infty. \tag{0.2}$$

In [BF91, Theorem 1.1] Bożejko and Fendler shows that, for countable discrete groups, all Herz–Schur multipliers can be realized as coefficients of representations satisfying a condition similar to (0.2). More specifically, they show that if φ is a Hermitian Herz–Schur multiplier on a countable discrete group Γ, then there exists a representation $(\pi, \mathscr{H})$ and vectors $\xi, \eta \in \mathscr{H}$ such that

$$\varphi(y^{-1}x) = \langle \pi(x)\xi, \pi(y)\eta \rangle \qquad (x, y \in \Gamma),$$

with

$$\sup_{x \in \Gamma} \|\pi(x)\xi\| \leq \|\varphi\|_{M_0A(\Gamma)}^{\frac{1}{2}} \quad \text{and} \quad \sup_{y \in \Gamma} \|\pi(y)\eta\| \leq \|\varphi\|_{M_0A(\Gamma)}^{\frac{1}{2}}.$$

Furthermore, they give a quantitative bound on $\|\pi(g)\|$ for $g \in \Gamma$ and note that the same holds for non-Hermitian Herz–Schur multipliers by including a square root two factor in the bound for $\sup_{x\in\Gamma}\|\pi(x)\xi\|$ and $\sup_{y\in\Gamma}\|\pi(y)\eta\|$. In section 1 we present a generalization of the result by Bożejko and Fendler to second countable, locally compact groups (cf. Theorem 1.1 and Corollary 1.4). We also present the following theorem based on criterion (0.2) (cf. Theorem 1.5):

0.2 Theorem. *Let G be a second countable, locally compact group and let* d *be a proper, left invariant metric on G which has* at most exponential growth*, i.e.,*

$$\mu(B_n(e)) \leq a \cdot e^{bn} \qquad (n \in \mathbb{N})$$

for some constants $a, b > 0$, where μ is a left invariant Haar measure on G and $B_n(e) = \{g \in G \,:\, \mathrm{d}(g, e) < n\}$ is the open ball of radius n centered around the identity element $e \in G$. Then for any continuous Herz–Schur multiplier φ on G there exists a strongly continuous representation $(\pi, \mathscr{H})$ and vectors $\xi, \eta \in \mathscr{H}$ such that

$$\varphi(y^{-1}x) = \langle \pi(x)\xi, \pi(y^{-1})^*\eta \rangle \qquad (x, y \in G),$$

with

$$\sup_{x\in G}\|\pi(x)\xi\| = \|\varphi\|_{M_0A(G)}^{\frac{1}{2}} \quad \textit{and} \quad \sup_{y\in G}\|\pi(y^{-1})^*\eta\| = \|\varphi\|_{M_0A(G)}^{\frac{1}{2}}.$$

Moreover, for every fixed $c > b$, $(\pi, \mathscr{H})$ can be chosen such that

$$\|\pi(g)\| \leq e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

Note that the existence of a proper, left invariant metric with at most exponential growth on a second countable, locally compact group is guarantied by [HP06].

1 Coefficients of non-uniformly bounded representations

Second countability guarantees the existence of a proper, left invariant metric, cf. [Str74, Theorem]. Actually, according to Haagerup and Przybyszewska (cf. [HP06]) one can choose this metric, d, to have *at most exponential growth*, i.e.,

$$\mu(B_n(e)) \leq a \cdot e^{bn} \qquad (n \in \mathbb{N}) \tag{1.1}$$

for some constants $a, b > 0$, where μ is a left invariant Haar measure on G and $B_n(e) = \{g \in G \ : \ \mathrm{d}(g,e) < n\}$ is the open ball of radius n centered around the identity element $e \in G$.

Inspired by the proof of [BF91, Theorem 1.1], we state and prove Theorem 1.1 for *Hermitian* Herz–Schur multipliers, i.e., Herz–Schur multipliers φ for which $\varphi^* = \varphi$, where

$$\varphi^*(g) = \overline{\varphi(g^{-1})} \qquad (g \in G).$$

The general (not necessarily Hermitian) case is treated in Corollary 1.4 and Theorem 1.5.

1.1 Theorem. *If φ is a continuous Hermitian Herz–Schur multiplier on a second countable, locally compact group G, and* d *is a proper, left invariant metric on G satisfying* (1.1) *for some $a, b > 0$ (which exists according to [HP06]), then there exists a strongly continuous representation $(\pi, \mathscr{H})$ and vectors $\xi, \eta \in \mathscr{H}$ such that*

$$\varphi(y^{-1}x) = \langle \pi(x)\xi, \pi(y)\eta \rangle \qquad (x, y \in G),$$

with

$$\sup_{x \in G} \|\pi(x)\xi\| = \|\varphi\|_{M_0A(G)}^{\frac{1}{2}} \quad \textit{and} \quad \sup_{y \in G} \|\pi(y)\eta\| = \|\varphi\|_{M_0A(G)}^{\frac{1}{2}}.$$

Moreover, for every fixed $c > b$, $(\pi, \mathscr{H})$ can be chosen such that

$$\|\pi(g)\| \leq \ e^{\frac{c}{2} \cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

Before we proceed with the proof of Theorem 1.1 we need the following application of the above mentioned result from [HP06], which was communicated to us by Haagerup.

1.2 Lemma. *If G is a second countable, locally compact group, then there exists a positive function $h \in L^1(G)$ with $\|h\|_1 = 1$, and a positive function c on G such that*

$$\frac{1}{c(g)} \int_G f(z)h(z) \mathrm{d}\mu(z) \leq \int_G f(z)h(gz)\mathrm{d}\mu(z) \leq c(g) \int_G f(z)h(z)\mathrm{d}\mu(z)$$

for $g \in G$ and any positive $f \in L^\infty(G)$, where μ is the Haar measure on G. Moreover, we may use

$$h(g) = \frac{e^{-c \cdot \mathrm{d}(g,e)}}{\int_G e^{-c \cdot \mathrm{d}(x,e)} \mathrm{d}\mu(x)} \quad \textit{and} \quad c(g) = e^{c \cdot \mathrm{d}(g,e)} \qquad (g \in G)$$

for $c > b$ when d *is a proper, left invariant metric on G satisfying* (1.1).

Proof. Let μ be a left invariant Haar measure on G and let d be a proper, left invariant metric on G satisfying (1.1). We claim that

$$0 < \int_G e^{-c\cdot \mathrm{d}(g,e)} \mathrm{d}\mu(g) < \infty.$$

Put $E_1 = B_1(e)$ and define inductively

$$E_n = B_n(e) \setminus B_{n-1}(e) \qquad (n \geq 2).$$

Then $G = \bigsqcup_{n=1}^{\infty} E_n$ and

$$e^{-cn} \leq e^{-c\cdot \mathrm{d}(g,e)} \leq e^{-c(n-1)} \qquad (g \in E_n).$$

Hence,

$$\begin{aligned} \int_G e^{-c\cdot \mathrm{d}(g,e)} \mathrm{d}\mu(g) &= \sum_{n=1}^{\infty} \int_{E_n} e^{-c\cdot \mathrm{d}(g,e)} \mathrm{d}\mu(g) \\ &\leq \sum_{n=1}^{\infty} e^{-c(n-1)} \mu(E_n) \\ &\leq e^c \sum_{n=1}^{\infty} e^{-cn} \mu(B_n(e)) \\ &\leq a e^c \sum_{n=1}^{\infty} e^{(b-c)n} \\ &< \infty \end{aligned}$$

because $c > b$.

By the reverse triangle inequality we see that

$$|\mathrm{d}(z, g^{-1}) - \mathrm{d}(z, e)| \leq \mathrm{d}(e, g^{-1}) \qquad (g, z \in G).$$

Using left invariance of the metric one finds that

$$|\mathrm{d}(gz, e) - \mathrm{d}(z, e)| \leq \mathrm{d}(g, e) \qquad (g, z \in G).$$

This implies

$$\frac{1}{c(g)} e^{-c\cdot \mathrm{d}(z,e)} \leq e^{-c\cdot \mathrm{d}(gz,e)} \leq c(g) e^{-c\cdot \mathrm{d}(z,e)} \qquad (g, z \in G),$$

which is easily seen to complete the proof. ☐

1.3 Lemma. *Assume that $\mathscr{H}$ is a Hilbert space and that $R : G \to \mathscr{H}$ is bounded and continuous. Let $R' : G \to L^2(G, \mathscr{H}, \mu)$ be given by*

$$R'(x)(z) = \sqrt{h(z)}R(z^{-1}x) \qquad (z \in G)$$

for $x \in G$, then R' is bounded and continuous, with $\|R'(x)\|_2 \le \|R\|_\infty$ for all $x \in G$. Also, let $\mathscr{K}_R = \overline{\operatorname{span}}\{R'(x) \, : \, x \in G\}$ be a sub-Hilbert space of $L^2(G, \mathscr{H}, \mu)$. Then there exists a unique representation $(\pi_R, \mathscr{K}_R)$ such that

$$\pi_R(g)R'(x) = R'(gx) \qquad (g, x \in G).$$

Moreover,

$$\|\pi_R(g)\| \le e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G)$$

and the representation is strongly continuous.

Proof. From Lebesgue's dominated convergence theorem it easily follows that R' is continuous. To see that R' is bounded, note that

$$\|R'(x)\|_2^2 = \int_G h(z)\|R(z^{-1}x)\|^2 \mathrm{d}\mu(z) \le \|R\|_\infty^2 \qquad (x \in G).$$

If $n \in \mathbb{N}$, $x_1, \ldots, x_n \in G$ and $c_1, \ldots, c_n \in \mathbb{C}$, then Lemma 1.2 implies that

$$\int_G \|\sum_{i=1}^n c_i R(z^{-1}x_i)\|^2 h(gz)\mathrm{d}\mu(z) \le c(g) \int_G \|\sum_{i=1}^n c_i R(z^{-1}x_i)\|^2 h(z)\mathrm{d}\mu(z)$$

for $g \in G$, where

$$c(g) = e^{c\cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

It follows that

$$\|\sum_{i=1}^n c_i R'(gx_i)\|_2^2 \le c(g)\|\sum_{i=1}^n c_i R'(x_i)\|_2^2 \qquad (g \in G),$$

from which we conclude that there exists a unique representation $(\pi_R, \mathscr{K}_R)$ of G such that

$$\pi_R(g)R'(x) = R'(gx) \qquad (g, x \in G).$$

Furthermore,

$$\|\pi_R(g)\| \le \sqrt{c(g)} \qquad (g \in G).$$

We proceed to show that the representation is strongly continuous. Since $\operatorname{span}\{R'(x) \, : \, x \in G\}$ is total in $\mathscr{K}_R$ and $\|\pi_R(g)\| \le \sqrt{c(g)}$, where $g \mapsto \sqrt{c(g)}$ is a continuous function, it is enough to show that

$$\lim_{n\to\infty} \pi_R(g_n)R'(x) = R'(x)$$

for $x \in G$, when $(g_n)_{n\in\mathbb{N}}$ is a sequence converging to the identity $e \in G$ (since G is second countable we do not have to consider nets). But $\pi_R(g_n)R'(x) = R'(g_n x)$ so this follows simply from continuity of R'. □

Proof of Theorem 1.1. Assume that φ is a continuous Hermitian Herz–Schur multiplier and use Proposition 0.1 to find a Hilbert space $\mathscr{H}$ and bounded, continuous maps $P, Q : G \to \mathscr{H}$ such that

$$\varphi(y^{-1}x) = \langle P(x), Q(y)\rangle \qquad (x, y \in G)$$

and

$$\|P\|_\infty = \|Q\|_\infty = \|\varphi\|_{M_0A(G)}^{\frac{1}{2}}.$$

Define

$$a_\pm(x, y) = \frac{1}{4}\langle P(x) \pm Q(x), P(y) \pm Q(y)\rangle \qquad (x, y \in G).$$

This gives rise to two positive definite, bounded kernels on $G \times G$ satisfying

$$a_+(x, y) - a_-(x, y) = \frac{1}{2}\varphi(y^{-1}x) + \frac{1}{2}\varphi^*(y^{-1}x) = \varphi(y^{-1}x) \qquad (x, y \in G)$$

and

$$a_+(x, x) + a_-(x, x) = \frac{1}{2}\|P(x)\|^2 + \frac{1}{2}\|Q(x)\|^2 \le \|\varphi\|_{M_0A(G)} \qquad (x \in G).$$

Let

$$h(g) = \frac{e^{-c\cdot \mathrm{d}(g,e)}}{\int_G e^{-c\cdot \mathrm{d}(x,e)}\mathrm{d}\mu(x)} \qquad (g \in G)$$

for some $c > b$, when d is a proper, left invariant metric on G satisfying (1.1) (cf. Lemma 1.2). Define $(P \pm Q)' : G \to L^2(G, \mathscr{H}, \mu)$ by

$$(P \pm Q)'(x)(z) = \sqrt{h(z)}(P \pm Q)(z^{-1}x) \qquad (z \in G)$$

for $x \in G$. According to Lemma 1.3 there exist strongly continuous representations $(\pi_{P\pm Q}, \mathscr{K}_{P\pm Q})$, where $\mathscr{K}_{P\pm Q} = \overline{\mathrm{span}}\{(P' \pm Q')(x) \; : \; x \in G\}$ and $\pi_{P\pm Q}(g)(P \pm Q)'(x) = (P \pm Q)'(gx)$ for $g, x \in G$. Furthermore, these representations satisfy

$$\|\pi_{P\pm Q}(g)\| \le e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G). \tag{1.2}$$

Put

$$A_\pm(x, y) = \langle (P \pm Q)'(x), (P \pm Q)'(y)\rangle_{\mathscr{K}_{P\pm Q}} \qquad (x, y \in G).$$

Then $A_\pm$ are positive definite, bounded kernels on $G \times G$ satisfying

$$A_+(x, y) - A_-(x, y) = \varphi(y^{-1}x) \qquad (x, y \in G) \tag{1.3}$$

and

$$A_+(x,x) + A_-(x,x) \le \|\varphi\|_{M_0A(G)} \qquad (x \in G). \tag{1.4}$$

To make the notation less cumbersome, let $\pi_\pm = \pi_{P\pm Q}$ and $\mathscr{K}_\pm = \mathscr{K}_{P\pm Q}$ and define $\xi_\pm = (P \pm Q)'(e)$. Notice that

$$\langle \pi_\pm(x)\xi_\pm, \pi_\pm(y)\xi_\pm \rangle_{\mathscr{K}_\pm} = A_\pm(x,y) \qquad (x,y \in G),$$

and that (1.2) now reads

$$\|\pi_\pm(g)\| \le e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

Put

$$\mathscr{K} = \mathscr{K}_+ \oplus \mathscr{K}_-, \quad \xi = \xi_+ \oplus \xi_-, \quad \eta = \xi_+ \oplus -\xi_- \quad \text{and} \quad \pi = \pi_+ \oplus \pi_-.$$

Observe that π is a strongly continuous representation such that

$$\|\pi(g)\| \le e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G) \tag{1.5}$$

and

$$\langle \pi(x)\xi, \pi(y)\eta \rangle_{\mathscr{K}} = \varphi(y^{-1}x) \qquad (x,y \in G).$$

Finally, observe that

$$\|\pi(x)\xi\|^2 = \|\pi_+(x)\xi_+\|^2 + \|\pi_-(x)\xi_-\|^2 = A_+(x,x) + A_-(x,x) \le \|\varphi\|_{M_0A(G)}$$

for $x \in G$, and similarly

$$\|\pi(y)\eta\|^2 \le \|\varphi\|_{M_0A(G)}$$

for $y \in G$. This finishes the proof. ☐

1.4 Corollary. *If φ is a continuous Herz–Schur multiplier on a second countable, locally compact group G, and d is a proper, left invariant metric on G satisfying* (1.1) *for some $a, b > 0$ (which exists according to [HP06]), then there exists a strongly continuous representation $(\pi, \mathscr{H})$ and vectors $\xi, \eta \in \mathscr{H}$ such that*

$$\varphi(y^{-1}x) = \langle \pi(x)\xi, \pi(y)\eta \rangle \qquad (x,y \in G),$$

with

$$\sup_{x\in G} \|\pi(x)\xi\| \le \sqrt{2}\|\varphi\|_{M_0A(G)}^{\frac{1}{2}} \quad \textit{and} \quad \sup_{y \in G}\|\pi(y)\eta\| \le \sqrt{2}\|\varphi\|_{M_0A(G)}^{\frac{1}{2}}.$$

Moreover, for every fixed $c > b$, $(\pi, \mathscr{H})$ can be chosen such that

$$\|\pi(g)\| \le e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

Proof. This follows from Theorem 1.1 since

$$\varphi = \mathrm{Re}(\varphi) + i\mathrm{Im}(\varphi),$$

where

$$\mathrm{Re}(\varphi) = \frac{\varphi + \varphi^*}{2} \quad \text{and} \quad \mathrm{Im}(\varphi) = \frac{\varphi - \varphi^*}{2i}$$

are continuous Hermitian Herz–Schur multipliers with

$$\|\mathrm{Re}(\varphi)\|_{M_0A(G)} \leq \|\varphi\|_{M_0A(G)} \quad \text{and} \quad \|\mathrm{Im}(\varphi)\|_{M_0A(G)} \leq \|\varphi\|_{M_0A(G)}.$$

□

1.5 Theorem. *If φ is a continuous Herz–Schur multiplier on a second countable, locally compact group G, and d is a proper, left invariant metric on G satisfying* (1.1) *for some $a, b > 0$ (which exists according to [HP06]), then there exists a strongly continuous representation $(\pi, \mathscr{H})$ and vectors $\xi, \eta \in \mathscr{H}$ such that*

$$\varphi(y^{-1}x) = \langle \pi(x)\xi, \pi(y^{-1})^*\eta \rangle \qquad (x, y \in G),$$

with

$$\sup_{x\in G} \|\pi(x)\xi\| = \|\varphi\|^{\frac{1}{2}}_{M_0A(G)} \quad \textit{and} \quad \sup_{y\in G} \|\pi(y^{-1})^*\eta\| = \|\varphi\|^{\frac{1}{2}}_{M_0A(G)}.$$

Moreover, for every fixed $c > b$, $(\pi, \mathscr{H})$ can be chosen such that

$$\|\pi(g)\| \leq e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

Proof. Assume that φ is a continuous Herz–Schur multiplier and use Proposition 0.1 to find a Hilbert space $\mathscr{H}$ and bounded, continuous maps $P, Q : G \to \mathscr{H}$ such that

$$\varphi(y^{-1}x) = \langle P(x), Q(y) \rangle \qquad (x, y \in G)$$

and

$$\|P\|_\infty = \|Q\|_\infty = \|\varphi\|^{\frac{1}{2}}_{M_0A(G)}.$$

Let

$$h(g) = \frac{e^{-c\cdot \mathrm{d}(g,e)}}{\int_G e^{-c\cdot \mathrm{d}(x,e)} \mathrm{d}\mu(x)} \qquad (g \in G)$$

for some $c > b$, when d is a proper, left invariant metric on G satisfying (1.1) (cf. Lemma 1.2). Define $P', Q' : G \to L^2(G, \mathscr{H}, \mu)$ by

$$P'(x)(z) = \sqrt{h(z)}P(z^{-1}x) \quad \text{and} \quad Q'(y)(z) = \sqrt{h(z)}Q(z^{-1}y) \qquad (z \in G)$$

for $x, y \in G$. According to Lemma 1.3 there exists a strongly continuous representation $(\pi_P, \mathscr{K}_P)$, where $\mathscr{K}_P = \overline{\operatorname{span}}\{P'(x) : x \in G\}$ and $\pi_P(g)P'(x) = P'(gx)$ for $g, x \in G$. Furthermore, this representation satisfies

$$\|\pi_P(g)\| \leq e^{\frac{c}{2}\cdot \mathrm{d}(g,e)} \qquad (g \in G).$$

Observe that

$$\|P'(x)\|_2^2, \|Q'(y)\|_2^2 \leq \|\varphi\|_{M_0A(G)}$$

and

$$\langle P'(x), Q'(y)\rangle_{L^2(G,\mathscr{H},\mu)} = \int_G h(z)\langle P(z^{-1}x), Q(z^{-1}y)\rangle_{\mathscr{H}} \mathrm{d}\mu(z) = \varphi(y^{-1}x)$$

for $x, y \in G$. Put $\xi = P'(e)$ and $\eta = P_{\mathscr{K}_P}Q'(e)$, where $P_{\mathscr{K}_P}$ is the orthogonal projection on $\mathscr{K}_P$. Note that $\xi, \eta \in \mathscr{K}_P$ and

$$\varphi(y^{-1}x) = \langle \pi_P(y^{-1}x)\xi, \eta\rangle_{\mathscr{K}_P} = \langle \pi_P(x)\xi, \pi_P(y^{-1})^*\eta\rangle_{\mathscr{K}_P} \qquad (x, y \in G).$$

It is clear that $\|\pi_P(x)\xi\|_{\mathscr{K}_P}^2 = \|P'(x)\|_2^2 \leq \|\varphi\|_{M_0A(G)}$. The corresponding result for $\|\pi_P(y^{-1})^*\eta\|_{\mathscr{K}_P}^2$ requires more work. For $x \in G$ arbitrary we find that

$$\begin{aligned}
\langle \pi_P(y^{-1})P'(x), P_{\mathscr{K}_P}Q'(e)\rangle_{\mathscr{K}_P} &= \langle P'(y^{-1}x), P_{\mathscr{K}_P}Q'(e)\rangle_{\mathscr{K}_P} \\
&= \langle P'(y^{-1}x), Q'(e)\rangle_{\mathscr{H}} \\
&= \varphi(y^{-1}x) \\
&= \langle P'(x), Q'(y)\rangle_{\mathscr{H}} \\
&= \langle P'(x), P_{\mathscr{K}_P}Q'(y)\rangle_{\mathscr{K}_P},
\end{aligned}$$

from which we conclude that $\pi_P(y^{-1})^* P_{\mathscr{K}_P}Q'(e) = P_{\mathscr{K}_P}Q'(y)$ and therefore

$$\|\pi_P(y^{-1})^*\eta\|_{\mathscr{K}_P}^2 = \|P_{\mathscr{K}_P}Q'(y)\|_{\mathscr{K}_P}^2 \leq \|Q'(y)\|_2^2 \leq \|\varphi\|_{M_0A(G)}.$$

□

1.6 Remark. For the free group on N generators ($2 \leq N < \infty$) the constants a, b in (1.1) may be chosen as $a = \frac{N}{(N-1)(2N-1)}$ and $b = \ln(2N-1)$. This implies that for $r > \sqrt{2N-1}$, the representations $(\pi, \mathscr{H})$ from Theorem 1.1, Corollary 1.4 and Theorem 1.5 may be chosen to satisfy $\|\pi(g)\| \leq r^{\mathrm{d}(g,e)}$ for all $g \in G$.

Acknowledgments

I would like to thank Ryszard Szwarc and Marek Bożejko for an inspiring stay in Wrocław during March 2009.

References

[BF84] Marek Bożejko and Gero Fendler. Herz–Schur multipliers and completely bounded multipliers of the Fourier algebra of a locally compact group. *Boll. Un. Mat. Ital. A (6)*, 3(2):297–302, 1984.

[BF91] Marek Bożejko and Gero Fendler. Herz–Schur multipliers and uniformly bounded representations of discrete groups. *Arch. Math. (Basel)*, 57(3):290–298, 1991.

[DCH85] Jean De Cannière and Uffe Haagerup. Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. *Amer. J. Math.*, 107(2):455–500, 1985.

[Eym64] Pierre Eymard. L'algèbre de Fourier d'un groupe localement compact. *Bull. Soc. Math. France*, 92:181–236, 1964.

[Her74] Carl Herz. Une généralisation de la notion de transformée de Fourier-Stieltjes. *Ann. Inst. Fourier (Grenoble)*, 24(3):xiii, 145–157, 1974.

[HP06] Uffe Haagerup and Agata Przybyszewska. Proper metrics on locally compact groups, and proper affine isometric actions on Banach spaces. 2006, `arXiv:math/0606794v1` [math.OA].

[Jol92] Paul Jolissaint. A characterization of completely bounded multipliers of Fourier algebras. *Colloq. Math.*, 63(2):311–313, 1992.

[Pis05] Gilles Pisier. Are unitarizable groups amenable? In *Infinite groups: geometric, combinatorial and dynamical aspects*, volume 248 of *Progr. Math.*, pages 323–362. Birkhäuser, Basel, 2005.

[Ste09] Troels Steenstrup. Fourier multiplier norms of spherical functions on the generalized Lorentz groups. 2009, `arXiv:0911.4977v1` [math.RT].

[Str74] Raimond A. Struble. Metrics in locally compact groups. *Compositio Math.*, 28:217–222, 1974.

Troels Steenstrup `<troelssj@imada.sdu.dk>`

Department of Mathematics and Computer Science, University of Southern Denmark, Campusvej 55, DK–5230 Odense M, Denmark.